

Medication Guide

TRUVADA® (tru-VAH-dah)
(emtricitabine and tenofovir disoproxil fumarate)
tablets

Read this Medication Guide before you start taking TRUVADA and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.

This Medication Guide provides information about two different ways that TRUVADA may be used (see the Medication Guide section “**What is TRUVADA?**” for important information about how TRUVADA may be used):

- to treat Human Immunodeficiency Virus-1 (HIV-1) infection, and
 - to reduce the risk of getting HIV-1 infection in adults who are HIV-negative
- HIV is the virus that causes AIDS (Acquired Immune Deficiency Syndrome).

What is the most important information I should know about TRUVADA?

If you also have hepatitis B virus (HBV) infection and take TRUVADA, your hepatitis B may become worse if you stop taking TRUVADA.

- Do not stop taking TRUVADA without first talking to your healthcare provider.
- Do not run out of TRUVADA. Refill your prescription or talk to your healthcare provider before your TRUVADA is all gone.
- If your healthcare provider stops TRUVADA, your healthcare provider will need to watch you closely for several months to check your hepatitis B infection, or give you a medication to treat hepatitis B.

Tell your healthcare provider about any new or unusual symptoms you may have after you stop taking TRUVADA.

For more information about side effects, see the section “**What are the possible side effects of TRUVADA?**” in this Medication Guide.

Other important information for people who take TRUVADA to help reduce their risk of getting HIV-1 infection:

Before taking TRUVADA to reduce your risk of getting HIV-1 infection:

- **You must be HIV-negative to start TRUVADA. You must get tested to make sure that you do not already have HIV-1 infection.**
- **Do not take TRUVADA to reduce the risk of getting HIV-1 unless you are confirmed to be HIV-negative.**
- Many HIV-1 tests can miss HIV-1 infection in a person who has recently become infected. If you have flu-like symptoms, you could have recently become infected with HIV-1. Tell your healthcare provider if you had a flu-like illness within the last month before starting TRUVADA or at any time while taking TRUVADA. Symptoms of new HIV-1 infection include:
 - tiredness
 - fever
 - joint or muscle aches
 - headache
 - sore throat
 - vomiting or diarrhea
 - rash
 - night sweats
 - enlarged lymph nodes in the neck or groin

While you are taking TRUVADA to reduce your risk of getting HIV-1:

- **Just taking TRUVADA may not keep you from getting HIV-1.**
- **You must continue using safer sex practices while you are taking TRUVADA to reduce your risk of getting HIV-1.**
- **You must stay HIV-negative to keep taking TRUVADA to reduce your risk of infection.**
 - Know your HIV-1 status and the HIV-1 status of your partners.
 - Get tested for HIV-1 at least every 3 months or when your healthcare provider tells you.
 - Get tested for other sexually transmitted infections such as syphilis and gonorrhea. These infections make it easier for HIV-1 to infect you.
 - If you think you were exposed to HIV-1, tell your healthcare provider right away. They may want to do more tests to be sure you are still HIV-negative.
 - Get information and support to help reduce risky sexual behavior.

- Have fewer sex partners.
- Do not miss any doses of TRUVADA. Missing doses may increase your risk of getting HIV-1 infection.
- If you do become HIV-positive, you need more medicine than TRUVADA alone to treat HIV-1. TRUVADA by itself is not a complete treatment for HIV-1.
 - If you have HIV-1 and take only TRUVADA, over time your HIV-1 may become harder to treat.

See the section “**What should I avoid while taking TRUVADA?**” and talk to your healthcare provider for more information about how to prevent HIV-1 infection.

What is TRUVADA?

TRUVADA contains the prescription medicines emtricitabine (EMTRIVA®) and tenofovir disoproxil fumarate (VIREAD®). TRUVADA is used:

- **to treat HIV-1 infection when used with other HIV-1 medicines in adults and children who weigh at least 37 pounds (at least 17 kg).**
- **to help** reduce the risk of getting HIV-1 infection when used with safer sex practices in:
 - HIV-negative men who have sex with men, who are at high risk of getting infected with HIV-1 through sex.
 - Male-female sex partners when one partner has HIV-1 infection and the other does not.

Use of TRUVADA to treat HIV-1 infection:

- When used with other HIV-1 medicines to treat HIV-1 infection, TRUVADA may help:
 - Reduce the amount of HIV-1 in your blood. This is called “viral load”.
 - Increase the number of CD4+ (T) cells in your blood that help fight off other infections.

Reducing the amount of HIV-1 and increasing the CD4+ (T) cells in your blood may help improve your immune system. This may reduce your risk of death or getting infections that can happen when your immune system is weak (opportunistic infections).

- **TRUVADA does not cure HIV-1 or AIDS.** If you have HIV-1 infection, you must keep taking HIV-1 medicines to control HIV-1 infection and decrease HIV-related illnesses.
- It is not known if TRUVADA is safe and effective in children with HIV-1 infection who weigh less than 37 pounds (less than 17 kg).

Use of TRUVADA to reduce the risk of HIV-1 infection:

- **When used with safer sex practices, TRUVADA may help** to reduce the risk of getting HIV-1 infection:
 - TRUVADA works better to reduce the risk of getting HIV-1 when the medicines are in your bloodstream before you are exposed to HIV-1.

Who should not take TRUVADA?

For people using TRUVADA to reduce the risk of getting HIV-1 infection:

TRUVADA can only help reduce your risk of getting HIV-1 **before** you are infected. **Do not take TRUVADA to help reduce your risk of getting HIV-1 if:**

- you already have HIV-1 infection. If you are HIV-positive, you need to take other medicines with TRUVADA to treat HIV-1. TRUVADA by itself is not a complete treatment for HIV-1.
- you do not know your HIV-1 infection status. You may already be HIV-positive. You need to take other HIV-1 medicines with TRUVADA to treat HIV-1.

What should I tell my healthcare provider before taking TRUVADA?

Tell your healthcare provider if you:

- have liver problems including hepatitis B virus infection
 - have kidney problems or receive kidney dialysis treatment
 - have bone problems
 - have any other medical conditions
 - are pregnant or plan to become pregnant. It is not known if TRUVADA can harm your unborn baby. If you are a female who is taking TRUVADA to reduce the risk of getting HIV-1 infection and you become pregnant while taking TRUVADA, talk to your healthcare provider to decide if you should keep taking TRUVADA.
- Pregnancy Registry:** A pregnancy registry collects information about your health and the health of your baby. There is a pregnancy registry for women who take medicines to treat or prevent HIV-1 during pregnancy. For more information about the registry and how it works, talk to your healthcare provider.
- are breastfeeding or plan to breastfeed.
 - You should not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby.

- Do not breastfeed if you take TRUVADA. TRUVADA can pass to your baby in your breast milk.
- Talk with your healthcare provider about the best way to feed your baby.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Do not take TRUVADA if you also take any of the medicines listed below:

- medicines which also contain emtricitabine or tenofovir disoproxil fumarate, (ATRIPLA[®], COMPLERA[®], EMTRIVA, GENVOYA[®], ODEFSEY[®], STRIBILD[®], VEMLIDY[®], or VIREAD). These medicines contain one or more of the same active ingredients as TRUVADA.
- medicines which contain tenofovir alafenamide (DESCOVY[®], GENVOYA[®], or ODEFSEY[®])
- medicines which contain lamivudine (Combivir, Dutrebis, Epivir, Epivir-HBV, Epzicom, Triumeq, or Trizivir)
- adefovir (HEPSERA[®])

TRUVADA may interact with other medicines. Especially tell your healthcare provider if you take:

- didanosine (Videx EC)
- atazanavir (Reyataz)
- ledipasvir with sofosbuvir (HARVONI[®])
- sofosbuvir with velpatasvir (EPCLUSA[®])
- darunavir (Prezista)
- lopinavir with ritonavir (Kaletra)

Your healthcare provider may need to check you more often or change your dose if you take any of these medicines and TRUVADA.

Know the medicines you take. Keep a list of them to show your healthcare provider or pharmacist when you get a new medicine.

How should I take TRUVADA?

- Take TRUVADA exactly as prescribed.
- Take TRUVADA by mouth, with or without food.
- Children who take TRUVADA are prescribed a lower strength tablet than adults.
 - **Children should swallow the tablet whole.** Tell your healthcare provider if your child cannot swallow the tablet whole, because they may need a different HIV-1 medicine.
 - Your healthcare provider will change the dose of TRUVADA as needed based on your child's weight.
- TRUVADA is usually taken 1 time each day. Take TRUVADA at the same time each day to keep TRUVADA blood levels constant.
 - If you have kidney problems, your healthcare provider may tell you to take TRUVADA less often.
- Do not miss any doses of TRUVADA. Missing a dose lowers the amount of medicine in your blood.
- If you miss a dose of TRUVADA, take it as soon as you remember that day. Do not take more than 1 dose of TRUVADA in a day. Do not take 2 doses at the same time to make up for a missed dose. Call your healthcare provider or pharmacist if you are not sure what to do.
- Do not change your dose or stop taking TRUVADA without first talking with your healthcare provider. Stay under a healthcare provider's care when taking TRUVADA.
- Refill your TRUVADA prescription before you run out of medicine.
- If you take too much TRUVADA, call your healthcare provider or go to the nearest hospital emergency room right away.
- If you take TRUVADA to treat HIV-1 infection, you need to take other HIV-1 medicines. Your healthcare provider will tell you what medicines to take and how to take them.
- If you take TRUVADA to reduce your risk of getting HIV-1:
 - you must also use other methods to reduce your risk of getting HIV-1. See the section "**What should I avoid while taking TRUVADA?**" in this Medication Guide.
 - **Take TRUVADA every day**, not just when you think you have been exposed to HIV-1.

What should I avoid while taking TRUVADA?

While taking TRUVADA, avoid doing things that increase your risk of getting HIV-1 or spreading HIV-1 to other people.

- See the section "**What is the most important information I should know about TRUVADA?**" at the beginning of this Medication Guide.
- Do not have any kind of sex without protection. Always practice safer sex by using a latex or polyurethane condom, to lower the chance of sexual contact with semen, vaginal fluids, or blood.
- Do not share personal items that can have blood or body fluids on them, such as toothbrushes and razor blades.

- Do not share or re-use needles or other injection equipment.

Ask your healthcare provider if you have any questions about how to prevent getting HIV-1 or spreading HIV-1 to other people.

What are the possible side effects of TRUVADA?

TRUVADA may cause serious side effects, including:

- **See “What is the most important information I should know about TRUVADA?”**
- **New or worse kidney problems, including kidney failure.** If you had kidney problems in the past or take another medicine that can cause kidney problems, your healthcare provider may do blood tests to check your kidneys before you start and while you are taking TRUVADA. Your healthcare provider may tell you to take TRUVADA less often, or to stop taking TRUVADA if you have kidney problems.
- **Too much lactic acid in your blood (lactic acidosis).** Too much lactic acid is a serious but rare medical emergency that can lead to death. Tell your healthcare provider right away if you get these symptoms: weakness or being more tired than usual, unusual muscle pain, being short of breath or fast breathing, stomach pain with nausea and vomiting, cold or blue hands and feet, feel dizzy or lightheaded, or a fast or abnormal heartbeat.
- **Severe liver problems.** In rare cases, severe liver problems can happen that can lead to death. Tell your healthcare provider right away if you get these symptoms: skin or the white part of your eyes turns yellow, dark “tea-colored” urine, light-colored stools, loss of appetite for several days or longer, nausea, or stomach-area pain.
- **Bone problems** can happen in some people who take TRUVADA. Bone problems include bone pain, or softening or thinning of bones, which may lead to fractures. Your healthcare provider may need to do tests to check your bones.
- **Changes in your immune system (Immune Reconstitution Syndrome)** can happen when **an HIV-1 infected person starts taking HIV-1 medicines.** Your immune system may get stronger, and can then cause you to develop inflammation in areas of your body where infections may have been hiding for a long time. This inflammation may cause you to have minor symptoms, such as fever, but inflammation can also lead to serious problems. Tell your healthcare provider right away if you start having any new symptoms after starting TRUVADA for treatment of HIV-1 infection.

The most common side effects of TRUVADA in people taking TRUVADA to treat HIV-1 infection include:

- diarrhea
- nausea
- tiredness
- headache
- dizziness
- depression
- problems sleeping
- abnormal dreams
- rash

Common side effects in people who take TRUVADA to reduce the risk of getting HIV-1 infection include:

- stomach-area (abdomen) pain
- headache
- decreased weight

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of TRUVADA. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store TRUVADA?

- Store TRUVADA at room temperature between 68 °F to 77 °F (20 °C to 25 °C).
- Keep TRUVADA in its original container and keep the container tightly closed.
- Do not use TRUVADA if seal over bottle opening is broken or missing.

Keep TRUVADA and all other medicines out of reach of children.

General information about TRUVADA.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use TRUVADA for a condition for which it was not prescribed. Do not give TRUVADA to other people, even if they have the same symptoms you have. It may harm them.

This Medication Guide summarizes the most important information about TRUVADA. If you would like more information, talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about TRUVADA that is written for health professionals. For more information, call 1-800-445-3235 or go to www.TRUVADA.com.

What are the ingredients in TRUVADA?

Active ingredients: emtricitabine and tenofovir disoproxil fumarate.

Inactive ingredients: Croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose, and

pregelatinized starch (gluten free). The 200 mg/300 mg strength tablets are coated with Opadry II Blue Y-30-10701, which contains FD&C Blue #2 aluminum lake, hypromellose 2910, lactose monohydrate, titanium dioxide, and triacetin. The 167 mg/250 mg, 133 mg/200 mg, and 100 mg/150 mg strength tablets are coated with Opadry II Blue, which contains FD&C Blue #2 aluminum lake, hypromellose 2910, lactose monohydrate, titanium dioxide, and triacetin.

Manufactured for and distributed by:

Gilead Sciences, Inc.

Foster City, CA 94404

COMPLERA, DESCOVY, EMTRIVA, EPCLUSA, HARVONI, HEPSERA, GENVOYA, ODEFSEY, STRIBILD, TRUVADA, VEMLIDY, and VIREAD are trademarks of Gilead Sciences, Inc., or its related companies. ATRIPLA is a trademark of Bristol-Myers Squibb & Gilead Sciences, LLC. All other trademarks referenced herein are the property of their respective owners.

© 2017 Gilead Sciences, Inc. All rights reserved.

21752-GS-031

This Medication Guide has been approved by the U.S. Food and Drug Administration

Revised: April 2017